



# Battle between plants as antioxidants with free radicals in human body

Fatemeh Jamshidi-Kia<sup>1</sup>, Joko Priyanto Wibowo<sup>2,3</sup>, Mostafa Elachouri<sup>4,5</sup>, Rohollah Masumi<sup>6</sup>, Alizamen Salehifard-Jouneghani<sup>7</sup>, Zohreh Abolhassanzadeh<sup>8</sup>, Zahra Lorigooini<sup>8\*</sup>

<sup>1</sup>Department of Horticulture, Faculty of Agriculture, Shahrekord University, Shahrekord, Iran

<sup>2</sup>Faculty of Pharmacy, University of Muhammadiyah Banjarmasin, Banjarmasin, Indonesia

<sup>3</sup>Chemical and Pharmaceutical Biology Department, Groningen Research Institute of Pharmacy, University of Groningen, The Netherlands

<sup>4</sup>Mohammed First University, Sciences Faculty, Laboratory of Physiology, Genetic and Ethnopharmacology, Oujda, Morocco

<sup>5</sup>Award CV Raman International Fellowship for African Research 2010

<sup>6</sup>Department of Internal Medicine, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>7</sup>Department of Pediatrics, Clinical Research Development Unit, Hajar Hospital, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>8</sup>Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

## ARTICLE INFO

**Article Type:**  
Review

### Article History:

Received: 9 December 2019

Accepted: 10 April 2020

### Keywords:

Free radicals  
Antioxidant  
Plant  
Human health  
Oxidative stress

## ABSTRACT

Free radicals are constructed by natural physiological activities in the human cells as well as in the environment. They may be produced as a result of diet, smoking, exercise, inflammation, exposure to sunlight, air pollutants, stress, alcohol and drugs. Imbalanced redox status may lead to cellular oxidative stress, which can damage the cells of the body, resulting in an incidence of various diseases. If the endogenous antioxidants do not stop the production of reactive metabolites, they will be needed to bring about a balance in redox status. Natural antioxidants, for example plants, play an important part in this context. This paper seeks to report the available evidence about oxidative stress and the application of plants as antioxidant agents to fight free radicals in the human body. For this purpose, to better understand oxidative stress, the principles of free radical production, the role of free radicals in diseases, antioxidant defense mechanisms, and the role of herbs and diet in oxidative stress are discussed.

### Implication for health policy/practice/research/medical education:

This review highlights the available evidence about oxidative stress and the application of plants as antioxidant agents to fight free radicals in the human body. Also, the role of free radicals in diseases, antioxidant defense mechanisms, and the role of herbs and diet in oxidative stress were discussed.

*Please cite this paper as:* Jamshidi-Kia F, Wibowo JP, Elachouri M, Masumi R, Salehi Jouneghani A, Abolhassanzadeh Z, et al. Battle between plants as antioxidants with free radicals in human body. J Herbmec Pharmacol. 2020;9(3):191-199. doi: 10.34172/jhp.2020.25.

## Introduction

The recent enhancement of science about free radicals and reactive oxygen species (ROS) has led to developments in medicine (1). Free radicals are chemical compounds with unpaired electron(s). The formation of free radicals occurs in three paths: (a) through the hemolytic cleavage of a covalent bond of a regular molecule; (b) through removal of an electron from a regular molecule; (c) through connection of an electron to a regular molecule (2). Free radicals can also be considered as molecular species that exist independently and contain an unpaired electron. The existence of a free radical brings about certain properties

that are common to the majority of radicals. Radicals are mainly unstable and highly reactive. They can accept an electron from or donate it to other molecules, and therefore act as oxidants or reductants (3).

Antioxidants and unpaired electrons have turned into common terms in modern debates on the factors that may lead to diseases (4). When excessive amounts of free radicals and oxidants are produced, they result in a condition, namely, oxidative stress. This condition represents a harmful process that may change the cells structures considerably and destroy lipids, lipoproteins, and DNA (5). Hydroxyl (OH·), superoxide (O<sub>2</sub><sup>-</sup>) nitric

\*Corresponding author: Zahra Lorigooini,

Email: gueini.z@skums.ac.ir, zahralorigooini@gmail.com

oxide (NO·), and peroxy (RO<sub>2</sub>·) are a number of free radicals that are important for living organisms. Although hypochlorous acid (HOCl), peroxyxynitrite (ONOO<sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), singlet oxygen <sup>1</sup>Δg (1O<sub>2</sub>), and ozone (O<sub>3</sub>) can induce free-radical reactions in living organisms, they are not classified as free radicals. ROS is often used to refer to not only OH·, RO<sub>2</sub>·, NO· and O<sub>2</sub><sup>-</sup> but also <sup>1</sup>O<sub>2</sub>, ONOO<sup>-</sup>, HOCl, O<sub>3</sub>, and H<sub>2</sub>O<sub>2</sub>. ROS may be produced due to ultraviolet (UV) radiation, smoking, and other exogenous factors (6,7).

Factors inducing free radicals production can be exogenous or endogenous. A variety of antioxidant defenses have been developed to fight the constant oxidant load, but damages due to free radicals are unavoidably exacerbated as age advances. Denham Harman proposed the “Theory of Aging” and stated that the free radical load is the cause of general aging. It can also be the reason of certain diseases associated with the elderly (8). This article is aimed to present a comprehensive review on the available evidence about oxidative stress and the application of plants as antioxidant agents to fight free radicals in the human body. In this regard, the principles of free radical production, the role of free radicals in diseases, antioxidant defense mechanisms, and the role of herbs and diet in oxidative stress are discussed.

### Free radicals and human diseases

During the last two decades, accumulating evidence has reported the implication of free radical-mediated processes in a wide variety of human diseases (Figure 1) (9,10). When oxygen molecules split into single atoms with unpaired electrons, they turn into unstable free radicals that try to find other molecules or atoms to bind to. If this situation continues, a process called oxidative stress begins. Although production of free radicals under the normal physiological condition is adjusted by various antioxidants, production of excess reactive oxygen species leads to oxidative stress (11,12). Oxidative stress leads to cell injury, genes mutation and carcinogenesis (13,14). The progression of cancer requires molecular and cellular changes that are interposed by a variety of exogenous and endogenous stimuli (15). Oxidative stress has also been investigated in diseases such as amyotrophic lateral sclerosis, Parkinson’s and Alzheimer’s diseases (16,17). Studies have shown that oxidative damage can harm neurons and cause memory loss (15). Also, experimental studies have confirmed the role of oxidative stress in some cardiovascular diseases such as ischemia, cardiomyopathy, congestive heart failure, atherosclerosis, hypertension and cardiac hypertrophy (18-20). Increasing experimental and clinical evidence suggests that oxidative stress can be the pathogenesis of diabetes. Increase of free radicals and the concomitant mitigation of antioxidants lead to increase in lipid peroxidation, development of insulin resistance and damage to cellular organelles and enzymes, which can




Figure 1. Oxidative stress and diseases (10). GSH: Glutathione.

result in diabetes mellitus complications (21,22).

### Antioxidant defence mechanism

Antioxidants are chemical compounds which contain monohydroxy/polyhydroxy phenol, which can reduce peroxidation (23). These antioxidants, based on their nature, and their source and mechanisms of action, can be classified into different groups (Figure 2).

Antioxidants have different mechanisms against free radicals, the catalytic systems that neutralize or divert ROS (Figure 3).


### Mechanism action of antioxidants

Antioxidants (AH) react with R, RO and ROO to generate RH, ROH, ROOH and a relatively unreactive phenoxyl radical (RH).

Antioxidants can neutralize free radicals by giving up electrons (Figure 4) (28).

### Synthetic antioxidants

Synthetic antioxidants are synthesized by chemical processes because they are not naturally formed. Synthetic antioxidants can be based on their mode of action, classified into different groups: primary and secondary synthetic antioxidants. The primary antioxidants hamper the production of free radicals in oxidation process (29,30). Secondary antioxidants act by decomposing hydroperoxides, which are formed in lipid oxidation process, into constant and compound (Figure 5) (31). Synthetic antioxidants have been shown to have both beneficial and harmful biological effects on the molecules,



**Figure 2.** Classification of antioxidants A) based on their nature; B) based on their source and C) based on their mechanisms of action (24, 25).

cells and organs. Because the carcinogenic effects of most synthetic antioxidants have been observed in high doses, their doses of use are recommended and controlled by certain legislating authorities, e.g. the FDA (32).

### Antioxidant properties for human health

The medicinal use of plants by humans may date back to at least 60 000 years (33,34). The plants are rich in natural products such as polyphenols, alkaloids, and terpenoids with various notable bioactivities, such as antimicrobial, anti-inflammatory, and antioxidants. Plants can also exert protective effects by inhibiting DNA oxidative damage. These nature-based substances can be found in any part of the plant or the plant by-products including fruit pomaces, seeds, peels, and pulps (35). Plant-based products are being extensively used as antioxidants


because they can protect cells and other organisms against oxidative damage (36).

Some plants have free radical scavenging molecules including phenolic compounds, vitamins, terpenoids, etc, which have antioxidant activity (37,38). Studies have proven antioxidants possess anti-inflammatory, anti-atherosclerotic, antitumor, antimutagenic, anti-carcinogenic and antibacterial activities (39).



There are numerous reports of various antioxidant properties among different herbal families. For example, some plants from different families with high level of antioxidant activity are presented in Table 1.

### Prevention of oxidative stress by diet

Dietary antioxidants can help other antioxidants inhibit oxidative stress. Nutrient antioxidant deficiency is a cause




**Figure 3.** Different mechanism of antioxidants against free radicals (26, 27).



**Figure 4.** Mechanism action of antioxidants, R (alkyl radical), RO (alkoxy radical), ROO (alkylperoxy radical), AH (antioxidants), A (Non radical products), ROOH (alkyl hydroperoxide).

of many degenerative and chronic diseases. Each nutrient has a unique structure and function of antioxidants, derived either directly or indirectly from the diet (62,63). Some studies have shown that low blood levels of antioxidants or low antioxidant intake increase the risk of various diseases. As a result, suitable antioxidant diet and nature-based antioxidant supplements as a constituent of wholesome lifestyles are being taken into-account with respect to protecting health against oxidative stress (25).

Diet can be a major source of antioxidants, many of which can adjust different processes that are associated with various diseases. Vegetables and fruits contain large amounts of antioxidants. These compounds can also be found in teas, nuts, legumes, grain cereals, teas, and other food products (64). Antioxidants that are absorbed by diet



**Figure 5.** Classification of antioxidants based on their mode of action (29,30).

contribute greatly to helping endogenous antioxidants neutralize excess free radicals. Vitamins, co-enzyme Q10, N-acetyl cysteine, alpha-lipoic acid, carotenoids, phenolic compounds, etc. are commonly used as nutritive and dietary antioxidants (64,65). The absorption of natural antioxidants from diet is, therefore, the first choice because natural antioxidants can play a considerable part in the prevention and adjunctive treatment of diseases (66).

#### Treatment of oxidative stress by diet

Numerous studies have shown that antioxidants available in foods, such as flavonoids, vitamin C and vitamin E serve as effective antioxidants in biological systems including lipoproteins, plasma, and cultured cells (67). For example, vitamin C efficiently inhibits the oxidation of lipid and protein in human plasma exposed to different (patho) physiologically relevant oxidative stress, e.g., reagent or myeloperoxidase-derived hypochlorous acid, activated polymorphonuclear leukocytes, redox-active iron or copper ions, and cigarette smoke (68). Vitamin E serves as a chain-breaking agent to deal with lipid peroxidation. Lycopene,  $\beta$ -carotene, lutein, and other carotenoids and oxy-carotenoids are efficient singlet oxygen quenchers and, therefore, may serve as important agents to defend the eye and other body organs against oxidative hurt induced by UV (69).

Many groups of plants have so far been used to cure diseases. Numerous drugs have been isolated from plant sources. For example, certain plant-derived alkaloids have been investigated for their potential use to treat Alzheimer's disease, and are being used for clinical purposes (e.g. galantamine derived from *Galanthus nivalis* L.) (70).

#### Controversial results of antioxidant in cell culture and in vivo studies

Cell culture is used to clarify the different mechanisms in the cell. Studies provide valuable information (71). Cell culture is often used to study the cellular effects of reactive species and of antioxidants (72). Oxygen ( $O_2$ ) concentrations range from 1–10 mm Hg in vivo. However,

**Table 1.** Plants with high level of antioxidant activity

Family	Species	Chemical Constituents	Reference
Asteraceae	<i>Achillea species</i>	Sesquiterpene lactone, polyenes, alkaloids, flavonoids, lignans and triterpenes	(40, 41)
Lauraceae	<i>Cinnamomum zeylanicum</i>	Eugenol and cinnamic aldehyde	(42)
Ericaceae	<i>Vaccinium oxycoccos</i>	Vitamin C and polyphenol	(43)
Fabaceae	<i>Melilotus officinalis</i>	Flavonoid and phenolic, hexadecanoic acid, lupanone, lupeol, betulinic acid, oleanolic and kaempferol-3-o- B-glu	(44)
Lamiaceae	<i>Lavandula officinalis</i>	Linalool, 1, 8 – cineole, borneol and camphor	(45, 46)
Lamiaceae	<i>Nepeta ispanhanica</i>	1,8-cineol, $\beta$ -pinene germacrene-D and $\alpha$ -pinene	(47)
Lamiaceae	<i>Salvia mirzayanii</i> Rech	Bicyclogermacrene, $\alpha$ - pinene, $\beta$ - pinene and sabinene	(48, 49)
Lamiaceae	<i>Ocimum sanctum</i> Linn	Eugenol, ursolic acid, estragole and thymol	(50)
Lamiaceae	<i>Thymus daenensis</i>	Thymol, gamma-terpinene and p-cymene	(51)
Leguminosae	<i>Glycyrrhiza glabra</i>	Glycyrrhizin, bitter principles, resins, mannite, and asparagine	(52)
Liliaceae	<i>Allium latifolium</i>	Thymol, dimethyl trisulfide, 4-methyl-5-thiazol ethanol, z- $\beta$ -cimene and carvacrol	(53)
Meliaceae	<i>Melia azedarach</i> L	Azardine, sterols, paraisine, rutin, palmitic, oleic acid and linoleic acid	(54)
Scrophulariaceae	<i>Verbascum songaricum</i>	Phenolic, 9-12 octadecadienoic acid, butanoic acid, 2,2-dimethyl butyl ester, heptyl ester propionic acid	(55, 56)
Theaceae	<i>Camellia sinensis</i>	Caffeine and theophylline	(57)
Verbenaceae	<i>Lantana camara</i> L	Theveside, theviridoside, geniposide, 8-epiloganin, lamiridoside, shanzhside methyl ester, triterpenoids, oleanolic acid, lantan olic acid lantanoses A and B	(58)
Zingiberaceae	<i>Curcuma longa</i>	Pigments curcumin, beta-pipene, camphene and Eugenol	(59, 60)
Zingiberaceae	<i>Zingiber officinale</i>	Terpenoid, volatile oil, shagoals, zingerone and peradols	(61)

cell culture experiments are performed at 95% atmospheric air/5% CO<sub>2</sub> an O<sub>2</sub> tension of approximately 150 mm Hg. Rates of production of ROS by cellular enzymes or by leakage from electron transport chains are O<sub>2</sub> at 10 mm Hg, and so, oxidative stress is due to generation of ROS in cells cultured at higher than this level. Moreover, the photochemical oxidation of polyphenolic compounds in the culture may be producing H<sub>2</sub>O<sub>2</sub> (73). In addition, cell culture is often deficient in antioxidants (74). Vitamin E and vitamin C are two important antioxidants but vitamin E is seldom added because it is insoluble in water, and vitamin C is not added because it is unstable, as a result, cells are deficient of these antioxidants. Studies have shown that antioxidants often have useful effects when added to cultured cells (75,76).

#### Negative results of antioxidants and possible mechanism

In many studies, antioxidants and pro-oxidants have been carefully studied and the results have shown that in some conditions, dietary antioxidants may behave as pro-oxidants (77,78). For example, Yen et al (79) studied the pro-oxidant properties of four flavonoids (quercetin, naringenin, hesperetin, and morin) in human lymphocytes. The production of superoxide anion radical and lipid peroxidation increased with increasing concentration of flavonoids. Many antioxidants may act, in some conditions, as pro-oxidants. Pro-oxidant process is directly proportional to the total number of hydroxyl groups in the flavonoids molecule (80-82). Therefore,


flavonoids cannot be considered merely antioxidants, since in their reactions they can also exhibit pro-oxidative activity. This activity may explain the observed toxicity of some flavonoids in vivo (83). However, this pro-oxidative behavior can also be useful, as it can increase the level of antioxidant defense and protein transfer enzymes and lead to overall protection in the body by imposing a mild degree of oxidative stress (84).

#### Relationship of structure-pro-oxidant activity to flavonoids


There is a connection between pro-oxidant ability and the presence of double bonds and ortho di/trihydroxylated B-ring phenol (Figure 6) (85).

The pro-oxidant behavior of flavonoids is dependent on their metal chelating ability, reducing properties and radical-scavenging ability. Also, flavonoids inhibit pro-oxidant enzymes. In the presence of O<sub>2</sub>, transition metals such as Fe or Cu catalyze the redox cycling of flavonoids, leading to the development of ROS (86). This may be related to the ability of flavonoids to commence autoxidation, catalyzed by transition metals, to yield superoxide anions, which would produce hydrogen peroxide and hydroxyl radicals (87). Flavonoids with A- or B-ring pyrogallol configurations induce DNA single-strand breakage (88). The process of DNA detriment takes place in such a way that catechol group in the ring A or B is oxidized by a Cu (II) ion bound to DNA which can damage generated ROS responsible for DNA damage





**Figure 6.** Relationship of structure-prooxidant activity to flavonoids.



**Figure 7.** The way of production of Fe (II)-flavonoid complex from Fe (III)-flavonoid complex (88).

(89,90). The oxidative DNA induced by Fe (III)-flavonoid complex (91), the way of production of Fe (II)-flavonoid complex from Fe (III)-flavonoid complex which then binds to DNA and generates ROS, is presented in [Figure 7](#).

### Conclusion

Recent medical developments indicate that some of the diseases are due to free radicals. The risk of oxidative stress-induced diseases is intensified by chemical exposure, unhealthy lifestyle, smoking, pollution, medications, stress, illness, etc. Antioxidants neutralize free radicals and decrease damage due to oxidative stress. Natural antioxidants and herbal drugs have been considered to be the main sources of antioxidant agents to prevent damage caused by free radicals. A diet containing various healthy foods with antioxidant properties is still the best strategy to benefit from antioxidants and the many other dietary bioactive components. However, in a few conditions antioxidants may act as pro-oxidant.

### Authors' contributions

ZL designed the research. All the authors contributed to data collection and preparation of the manuscript. The first draft was prepared by FJ. All authors read the final version and confirmed for the publication.

### Conflict of interests

The authors declared that there was no conflict of interest in the study.

### Ethical considerations

Ethical issues including text plagiarism, misconduct, manipulation or appropriation, data fabrication, falsification, redundant publication as well as duplicate submissions have been carefully observed by authors.

### Funding/Support

This work was supported by grant No. 1933 from Shahrekord University of Medical Sciences.

### References

1. Aruoma OI. Methodological considerations for characterizing potential antioxidant actions of bioactive components in plant foods. *Mutat Res.* 2003;523-524:9-20. doi: 10.1016/s0027-5107(02)00317-2.

2. Sala A, Recio M, Giner RM, Máñez S, Tournier H, Schinella G, et al. Anti-inflammatory and antioxidant properties of *Helichrysum italicum*. *J Pharm Pharmacol.* 2002;54(3):365-71. doi: 10.1211/0022357021778600.
3. Cheeseman KH, Slater TF. An introduction to free radical biochemistry. *Br Med Bull.* 1993;49(3):481-93. doi: 10.1093/oxfordjournals.bmb.a072625.
4. Aruoma OI. Nutrition and health aspects of free radicals and antioxidants. *Food Chem Toxicol.* 1994;32(7):671-83. doi: 10.1016/0278-6915(94)90011-6.
5. Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev.* 2007;87(1):315-424. doi: 10.1152/physrev.00029.2006.
6. Bhattacharyya A, Chattopadhyay R, Mitra S, Crowe SE. Oxidative stress: an essential factor in the pathogenesis of gastrointestinal mucosal diseases. *Physiol Rev.* 2014;94(2):329-54. doi: 10.1152/physrev.00040.2012.
7. Liu Z, Ren Z, Zhang J, Chuang CC, Kandaswamy E, Zhou T, et al. Role of ROS and nutritional antioxidants in human diseases. *Front Physiol.* 2018;9:477. doi: 10.3389/fphys.2018.00477.
8. Packer L. Oxidative stress, antioxidants, aging and disease. In: Cutler RG, Packer L, Bertram J, Mori A, eds. *Oxidative Stress and Aging*. Basel: Birkhäuser Basel; 1995. p. 1-14. doi: 10.1007/978-3-0348-7337-6\_1.
9. Southorn PA, Powis G. Free radicals in medicine. II. Involvement in human disease. *Mayo Clin Proc.* 1988;63(4):390-408. doi: 10.1016/s0025-6196(12)64862-9.
10. Aruoma OI. Free radicals, oxidative stress, and antioxidants in human health and disease. *J AM Oil Chem Soc.* 1998;75(2):199-212. doi: 10.1007/s11746-998-0032-9.
11. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol.* 2007;39(1):44-84. doi: 10.1016/j.biocel.2006.07.001.
12. Fang YZ, Yang S, Wu G. Free radicals, antioxidants, and nutrition. *Nutrition.* 2002;18(10):872-9. doi: 10.1016/s0899-9007(02)00916-4.
13. Fellow C, Division P. Cancer and oxidative stress. *J Jpn Med Assoc.* 2001;124(11):535-9.
14. Alkadi H, Ifeanyi OE. A Review on Free Radicals and Antioxidants. *Int J Curr Res Med Sci.* 2018;4(2):123-33. doi: 10.22192/ijcrms.2018.04.02.019.
15. Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact.* 2006;160(1):1-40. doi: 10.1016/j.cbi.2005.12.009.

16. Singh RP, Sharad S, Kapur S. Free radicals and oxidative stress in neurodegenerative diseases: relevance of dietary antioxidants. *J Indian Acad Clin Med.* 2004;5(3):218-25.
17. Cenini G, Lloret A, Cascella R. Oxidative stress in neurodegenerative diseases: from a mitochondrial point of view. *Oxid Med Cell Longev.* 2019;2019:2105607. doi: 10.1155/2019/2105607.
18. Christen Y. Oxidative stress and Alzheimer disease. *Am J Clin Nutr.* 2000;71(2):621S-9S. doi: 10.1093/ajcn/71.2.621s.
19. Ceriello A. Possible role of oxidative stress in the pathogenesis of hypertension. *Diabetes Care.* 2008;31 Suppl 2:S181-4. doi: 10.2337/dc08-s245.
20. Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, et al. Oxidative stress, aging, and diseases. *Clin Interv Aging.* 2018;13:757-72. doi: 10.2147/cia.s158513.
21. Chatterjee M, Saluja R, Kanneganti S, Chinta S, Dikshit M. Biochemical and molecular evaluation of neutrophil NOS in spontaneously hypertensive rats. *Cell Mol Biol (Noisy-le-grand).* 2007;53(1):84-93.
22. Lyons TJ, Jenkins AJ. Glycation, oxidation, and lipoxidation in the development of the complications of diabetes: a carbonyl stress hypothesis. *Diabetes Rev (Alex).* 1997;5(4):365-91.
23. Young IS, Tate S, Lightbody JH, McMaster D, Trimble ER. The effects of desferrioxamine and ascorbate on oxidative stress in the streptozotocin diabetic rat. *Free Radic Biol Med.* 1995;18(5):833-40. doi: 10.1016/0891-5849(94)00202-u.
24. German JB. Food processing and lipid oxidation. *Adv Exp Med Biol.* 1999;459:23-50. doi: 10.1007/978-1-4615-4853-9\_3.
25. Mironćzuk-Chodakowska I, Witkowska AM, Zujko ME. Endogenous non-enzymatic antioxidants in the human body. *Adv Med Sci.* 2018;63(1):68-78. doi: 10.1016/j.advms.2017.05.005.
26. Percival M. Antioxidants. Advanced Nutrition Publications, Inc; 1998. p. 1-4.
27. Benzie IF. Evolution of dietary antioxidants. *Comp Biochem Physiol A Mol Integr Physiol.* 2003;136(1):113-26. doi: 10.1016/s1095-6433(02)00368-9.
28. Bembem K, Sadana B. Effect of different cooking methods on the antioxidant components of Carrot. *Biosci Discov.* 2014;5(1):112-6.
29. Bose KS, Vyas P, Singh M. Plasma non-enzymatic antioxidants-vitamin C, E, beta-carotenes, reduced glutathione levels and total antioxidant activity in oral sub mucous fibrosis. *Eur Rev Med Pharmacol Sci.* 2012;16(4):530-2.
30. Kahl R. Synthetic antioxidants: biochemical actions and interference with radiation, toxic compounds, chemical mutagens and chemical carcinogens. *Toxicology.* 1984;33(3-4):185-228. doi: 10.1016/0300-483x(84)90038-6.
31. Bellés M, Alonso V, Roncalés P, Beltrán JA. Sulfite-free lamb burger meat: antimicrobial and antioxidant properties of green tea and carvacrol. *J Sci Food Agric.* 2019;99(1):464-72. doi: 10.1002/jsfa.9208.
32. Venkatesh R, Sood D. A review of the physiological implications of antioxidants in food. Bachelor of Science Interactive Qualifying Project. Worcester, Massachusetts, US: Worcester Polytechnic Institute; 2011.
33. Chen C, Pearson AM, Gray JI. Effects of synthetic antioxidants (BHA, BHT and PG) on the mutagenicity of IQ-like compounds. *Food Chem.* 1992;43(3):177-83. doi: 10.1016/0308-8146(92)90170-7.
34. Jamshidi-Kia F, Lorigooini Z, Amini-Khoei H. Medicinal plants: past history and future perspective. *J Herbmed Pharmacol.* 2018;7(1):1-7. doi: 10.15171/jhp.2018.01.
35. Yuan H, Ma Q, Ye L, Piao G. The traditional medicine and modern medicine from natural products. *Molecules.* 2016;21(5). doi: 10.3390/molecules21050559.
36. Abdel-Lateif KS, Eldeab HA, Maghrabi IA. The plant natural products: their antioxidants, free radical scavengers, DNA protection and antimicrobial activities. *J Bioprocess Biotech.* 2016;6(9):293. doi: 10.4172/2155-9821.1000293.
37. Gupta VK, Sharma SK. Plants as natural antioxidants. *Nat Prod Radiance.* 2006;5(4):326-34.
38. Zheng W, Wang SY. Antioxidant activity and phenolic compounds in selected herbs. *J Agric Food Chem.* 2001;49(11):5165-70. doi: 10.1021/jf010697n.
39. Cai Y, Sun M, Corke H. Antioxidant activity of betalains from plants of the Amaranthaceae. *J Agric Food Chem.* 2003;51(8):2288-94. doi: 10.1021/jf030045u.
40. Owen RW, Giacosa A, Hull WE, Haubner R, Spiegelhalder B, Bartsch H. The antioxidant/anticancer potential of phenolic compounds isolated from olive oil. *Eur J Cancer.* 2000;36(10):1235-47. doi: 10.1016/s0959-8049(00)00103-9.
41. Nickavar B, Kamalinejad M, Haj-Yahya M, Shafaghi B. Comparison of the free radical scavenging activity of six Iranian achillea. species. *Pharm Biol.* 2006;44(3):208-12. doi: 10.1080/13880200600686590.
42. Demirci F, Demirci B, Gurbuz I, Yesilada E. Characterization and biological activity of *Achillea teretifolia* Willd. and *A. nobilis* L. subsp. *neilreichii* (Kerner) Formanek essential oils. *Turk J Biol.* 2008;33(2):129-36. doi: 10.3906/biy-0808-1.
43. Carlsen MH, Halvorsen BL, Holte K, Bøhn SK, Dragland S, Sampson L, et al. The total antioxidant content of more than 3100 foods, beverages, spices, herbs and supplements used worldwide. *Nutr J.* 2010;9:3. doi: 10.1186/1475-2891-9-3.
44. Bean H, Schuler C, Leggett RE, Levin RM. Antioxidant levels of common fruits, vegetables, and juices versus protective activity against in vitro ischemia/reperfusion. *Int Urol Nephrol.* 2010;42(2):409-15. doi: 10.1007/s11255-009-9639-5.
45. Pourmorad F, Hosseinimehr SJ, Shahabimajd N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr J Biotechnol.* 2006;5(11):1142-5.
46. Afsharypour S Azarbajeny N. Chemical constituents of the flower essential oil of *Lavandula officinalis* Chaix. from Isfahan (Iran). *Iranian Journal of Pharmaceutical Sciences.* 2003;2(3):169-72.
47. Bouayed J, Piri K, Rammal H, Dicko A, Desor F, Younos C, et al. Comparative evaluation of the antioxidant potential of some Iranian medicinal plants. *Food Chem.* 2007;104(1):364-8. doi: 10.1016/j.foodchem.2006.11.069.
48. Farag RS, Badei AZMA, El Baroty GSA. Influence of thyme and clove essential oils on cottonseed oil oxidation. *J Am Oil Chem Soc.* 1989;66(6):800-4. doi: 10.1007/BF02653671.
49. Mirza M, Bahernik Z, Jamzad Z. Essential oil components of *Salvia mirzayani* Rech. *Iranian Journal of Medicinal and*

- Aromatic Plants Research. 2003;19(2):278-84.
50. Moein S, Farzami B, Khaghani S, Moein MR, Larijani BA. Antioxidant properties and protective effect on cell cytotoxicity of *Salvia mirzayani*. Pharm Biol. 2007;45(6):458-63. doi: 10.1080/13880200701389144.
  51. Triveni, Kumar K, Singh AK, Kumar R, Gupta V, Tripathi K. *Ocimum sanctum* Linn: a review on phytopharmacology therapeutic potential of Tulsi. Int J Pharm Phytopharm Res. 2013;3(2):148-51.
  52. Khanahmadi M, Rezazadeh S, Shahrezaei F, Taran M. Study on chemical composition of essential oil and anti-oxidant and anti microbial properties of *Artemisia haussknechtii*. Journal of Medicinal Plants. 2009;8(31):132-41.
  53. Parvaiz M, Hussain K, Khalid S, Hussain N, Iram N, Hussain Z, et al. A review: medicinal importance of *Glycyrrhiza glabra* L. (Fabaceae family). Glob J Pharmacol. 2014;8(1):8-13. doi: 10.5829/idosi.gjp.2014.8.1.81179.
  54. Khanahmadi M, Rezazadeh S, Shahrezaei F, Taran M, Rezaei M. Study on chemical composition of the essential oil, antimicrobial and antioxidant activities of *Allium latifolium* Gilib. (Liliaceae) extract. J Essent Oil Bear Plants. 2011;14(3):345-53. doi: 10.1080/0972060X.2011.10643944.
  55. Nahak G, Sahu RK. In vitro antioxidative activity of *Azadirachta indica* and *Melia azedarach* leaves by DPPH scavenging assay. Nat Sci Sleep. 2010;8(4):22-8.
  56. Safi Z, Saeidi K, Lorigooini Z, Shirmardi HA. Evaluation of total phenols and antioxidant activity of Mullein (*Verbascum songaricum*) ecotypes. Journal of Shahrekord University of Medical Sciences. 2016;17(6):68-75. [Persian].
  57. Karimian V, Vahabi MR, Fazilati M, Soleimani F. Chemical composition in two species of *Verbascum* collected from natural habitats, southern Iran. Journal of Herbal Drugs. 2013;4(3):127-32.
  58. Morteza-Semnani K, Saeedi M, Nozadi Z. Comparison of antioxidant activity of extract of Green tea to commercial antioxidants in 2% hydroquinone cream. Journal of Medicinal Plants. 2005;13(1):36-45. [Persian].
  59. Hussain H, Hussain J, Al-Harrasi A, Shinwari ZK. Chemistry of some species genus *Lantana*. Pak J Bot. 2011;43:51-62.
  60. Braga ME, Leal PF, Carvalho JE, Meireles MA. Comparison of yield, composition, and antioxidant activity of turmeric (*Curcuma longa* L.) extracts obtained using various techniques. J Agric Food Chem. 2003;51(22):6604-11. doi: 10.1021/jf0345550.
  61. Basnet P, Skalko-Basnet N. Curcumin: an anti-inflammatory molecule from a curry spice on the path to cancer treatment. Molecules. 2011;16(6):4567-98. doi: 10.3390/molecules16064567.
  62. Morakinyo AO, Oludare GO, Aderinto OT, Tasdup A. Antioxidant and free radical scavenging activities of aqueous and ethanol extracts of *Zingiber officinale*. Biol Med. 2011;3(5):25-30.
  63. Wilcox JK, Ash SL, Catignani GL. Antioxidants and prevention of chronic disease. Crit Rev Food Sci Nutr. 2004;44(4):275-95. doi: 10.1080/10408690490468489.
  64. Chikara S, Nagaprashantha LD, Singhal J, Horne D, Awasthi S, Singhal SS. Oxidative stress and dietary phytochemicals: role in cancer chemoprevention and treatment. Cancer Lett. 2018;413:122-34. doi: 10.1016/j.canlet.2017.11.002.
  65. FAO/WHO Expert Consultation Bangkok. Thailand Human Vitamin and Mineral Requirements. Rome: FAO; 2001. p. 303.
  66. Bouayed J, Bohn T. Exogenous antioxidants--Double-edged swords in cellular redox state: health beneficial effects at physiologic doses versus deleterious effects at high doses. Oxid Med Cell Longev. 2010;3(4):228-37. doi: 10.4161/oxim.3.4.12858.
  67. Li S, Chen G, Zhang C, Wu M, Wu S, Liu Q. Research progress of natural antioxidants in foods for the treatment of diseases. Food Science and Human Wellness. 2014;3(3-4):110-6. doi: 10.1016/j.fshw.2014.11.002.
  68. Frei B, England L, Ames BN. Ascorbate is an outstanding antioxidant in human blood plasma. Proc Natl Acad Sci U S A. 1989;86(16):6377-81. doi: 10.1073/pnas.86.16.6377.
  69. Carr AC, Frei B. Human neutrophils oxidize low-density lipoprotein by a hypochlorous acid-dependent mechanism: the role of vitamin C. Biol Chem. 2002;383(3-4):627-36. doi: 10.1515/bc.2002.065.
  70. Mayne ST. Antioxidant nutrients and chronic disease: use of biomarkers of exposure and oxidative stress status in epidemiologic research. J Nutr. 2003;133 Suppl 3:933S-40S. doi: 10.1093/jn/133.3.933S.
  71. Howes MJ, Perry NS, Houghton PJ. Plants with traditional uses and activities, relevant to the management of Alzheimer's disease and other cognitive disorders. Phytother Res. 2003;17(1):1-18. doi: 10.1002/ptr.1280.
  72. Halliwell B. Cell culture, oxidative stress, and antioxidants: avoiding pitfalls. Biomed J. 2014;37(3):99-105. doi: 10.4103/2319-4170.128725.
  73. Halliwell B. Oxidative stress and cancer: have we moved forward? Biochem J. 2007;401(1):1-11. doi: 10.1042/bj20061131.
  74. Halliwell B. Are polyphenols antioxidants or pro-oxidants? what do we learn from cell culture and in vivo studies? Arch Biochem Biophys. 2008;476(2):107-12. doi: 10.1016/j.abb.2008.01.028.
  75. Kelley EE, Buettner GR, Burns CP. Relative alpha-tocopherol deficiency in cultured cells: free radical-mediated lipid peroxidation, lipid oxidizability, and cellular polyunsaturated fatty acid content. Arch Biochem Biophys. 1995;319(1):102-9. doi: 10.1006/abbi.1995.1271.
  76. Karlenius TC, Shah F, Yu WC, Hawkes HJ, Tinggi U, Clarke FM, et al. The selenium content of cell culture serum influences redox-regulated gene expression. Biotechniques. 2011;50(5):295-301. doi: 10.2144/000113666.
  77. Papp LV, Lu J, Holmgren A, Khanna KK. From selenium to selenoproteins: synthesis, identity, and their role in human health. Antioxid Redox Signal. 2007;9(7):775-806. doi: 10.1089/ars.2007.1528.
  78. Halliwell B. The antioxidant paradox: less paradoxical now? Br J Clin Pharmacol. 2013;75(3):637-44. doi: 10.1111/j.1365-2125.2012.04272.x.
  79. Yen GC, Duh PD, Tsai HL, Huang SL. Pro-oxidative properties of flavonoids in human lymphocytes. Biosci Biotechnol Biochem. 2003;67(6):1215-22. doi: 10.1271/bbb.67.1215.
  80. Villanueva C, Kross RD. Antioxidant-induced stress. Int J Mol Sci. 2012;13(2):2091-109. doi: 10.3390/ijms13022091.
  81. Cao G, Sofic E, Prior RL. Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships.



- Free Radic Biol Med. 1997;22(5):749-60. doi: 10.1016/s0891-5849(96)00351-6.
82. Galati G, Sabzevari O, Wilson JX, O'Brien PJ. Prooxidant activity and cellular effects of the phenoxyl radicals of dietary flavonoids and other polyphenolics. *Toxicology*. 2002;177(1):91-104. doi: 10.1016/s0300-483x(02)00198-1.
  83. Goldman R, Claycamp GH, Sweetland MA, Sedlov AV, Tyurin VA, Kisin ER, et al. Myeloperoxidase-catalyzed redox-cycling of phenol promotes lipid peroxidation and thiol oxidation in HL-60 cells. *Free Radic Biol Med*. 1999;27(9-10):1050-63. doi: 10.1016/S0891-5849(99)00140-9.
  84. Kessler M, Ubeaud G, Jung L. Anti- and pro-oxidant activity of rutin and quercetin derivatives. *J Pharm Pharmacol*. 2003;55(1):131-42. doi: 10.1211/002235702559.
  85. Eghbaliferiz S, Iranshahi M. Prooxidant activity of polyphenols, flavonoids, anthocyanins and carotenoids: updated review of mechanisms and catalyzing metals. *Phytother Res*. 2016;30(9):1379-91. doi: 10.1002/ptr.5643.
  86. Li Y, Trush MA. Reactive oxygen-dependent DNA damage resulting from the oxidation of phenolic compounds by a copper-redox cycle mechanism. *Cancer Res*. 1994;54(7 Suppl):1895s-8s.
  87. Canada AT, Giannella E, Nguyen TD, Mason RP. The production of reactive oxygen species by dietary flavonols. *Free Radic Biol Med*. 1990;9(5):441-9. doi: 10.1016/0891-5849(90)90022-b.
  88. Ohshima H, Yoshie Y, Auriol S, Gilibert I. Antioxidant and pro-oxidant actions of flavonoids: effects on DNA damage induced by nitric oxide, peroxynitrite and nitroxyl anion. *Free Radic Biol Med*. 1998;25(9):1057-65. doi: 10.1016/s0891-5849(98)00141-5.
  89. Yamashita N, Tanemura H, Kawanishi S. Mechanism of oxidative DNA damage induced by quercetin in the presence of Cu(II). *Mutat Res*. 1999;425(1):107-15. doi: 10.1016/s0027-5107(99)00029-9.
  90. Amić D, Davidović-Amić D, Beslo D, Rastija V, Lucić B, Trinajstić N. SAR and QSAR of the antioxidant activity of flavonoids. *Curr Med Chem*. 2007;14(7):827-45. doi: 10.2174/092986707780090954.
  91. El Amrani FB, Perelló L, Real JA, González-Alvarez M, Alzuet G, Borrás J, García-Granda S, et al. Oxidative DNA cleavage induced by an iron(III) flavonoid complex: synthesis, crystal structure and characterization of chlorobis(flavonolato)(methanol) iron(III) complex. *J Inorg Biochem*. 2006;100(7):1208-18. doi: 10.1016/j.jinorgbio.2006.01.036.